

Learning Activity 2:
Diagnostic approaches and tools for predicting clinical response

1. Dubberke ER, Han Z, Bobo L, et al. Impact of clinical symptoms on interpretation of diagnostic assays for *Clostridium difficile* infections. *J Clin Microbiol.* 2011;49:2887-2893.
(<http://www.ncbi.nlm.nih.gov/pubmed/21697328>)
2. Belmares J, Gerding DN, Tillotson G, Johnson S. Measuring the severity of *Clostridium difficile* infection: implications for management and drug development. *Expert Rev Anti Infect Ther.* 2008;6:897-908.
(<http://www.ncbi.nlm.nih.gov/pubmed/19053902>)
3. Cohen SH, Gerding DN, Johnson S, et al. Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the Society for Healthcare Epidemiology of America (SHEA) and the Infectious Diseases Society of America (IDSA). *Infect Control Hosp Epidemiol.* 2010;31:431-455.
(<http://www.ncbi.nlm.nih.gov/pubmed/20307191>)
4. Bauer MP, Kuijper EJ, van Dissel JT, et al. European Society of Clinical Microbiology and Infectious Diseases (ESCMID): treatment guidance document for *Clostridium difficile* infection (CDI). *Clin Microbiol Infect.* 2009;15:1067-1079. (<http://www.ncbi.nlm.nih.gov/pubmed/19929973>)
5. Fujitani S, George WL, Murthy AR. Comparison of clinical severity score indices for *Clostridium difficile* infection. *Infect Control Hosp Epidemiol.* 2011;32:220-8. (<http://www.ncbi.nlm.nih.gov/pubmed/21460506>)
6. Jardin CG, Palmer HR, Shah DN, et al. Assessment of treatment patterns and patient outcomes before vs after implementation of a severity-based *Clostridium difficile* infection treatment policy. *J Hosp Infect.* 2013.
(<http://www.ncbi.nlm.nih.gov/pubmed/23834988>)
7. Zar FA, Bakkanagari SR, Moorthi KM, Davis MB. A comparison of vancomycin and metronidazole for the treatment of *Clostridium difficile*-associated diarrhea, stratified by disease severity. *Clin Infect Dis.* 2007;45:302-307. (<http://www.ncbi.nlm.nih.gov/pubmed/17599306>)
8. Shah DN, Bhatt NS, Welch JK, Koo HL, Garey KW. Defining acute renal dysfunction as a criterion for the severity of *Clostridium difficile* infection in patients with community-onset vs hospital-onset infection. *J Hosp Infect.* 2013;83:294-9. (<http://www.ncbi.nlm.nih.gov/pubmed/23433867>)
9. Arora V, Kachroo S, Ghantoli SS, Dupont HL, Garey KW. High Horn's index score predicts poor outcomes in patients with *Clostridium difficile* infection. *J Hosp Infect.* 2011;79:23-6.
(<http://www.ncbi.nlm.nih.gov/pubmed/21700363>)